

On the C_λ -extended w_∞ -symmetry

J. Douari¹

Institut des Hautes Etudes Scientifiques, Le Bois-Marie, 35, route de Chartres, F-91440
Bures-sur-Yvette, France.

E. H. El Kinani²

The Abdus Salam International Center for Theoretical Physics, Strada Costiera 11,
34014, Trieste, Italy.

Abstract

Starting from the C_λ -extended oscillator algebras, we obtain a new deformed w_∞ -algebra. More precisely, we show that the C_λ -extended w_∞ -algebra generators may be expressed via the annihilation and creation operators of the C_λ -extended oscillator algebras a and $a^\dagger$ as an infinite-dimensional extension of the realization of $sp(2)$ algebra.

1 Introduction

Over the past decade, much attention has been paid to the oscillator algebras in connection with the important role investigated in many physical systems, and they provide an useful tool in the theory of Lie algebra representations. On the other hand, their deformations and extensions have given new mathematical objects called Quantum Groups or Quantum Algebras [1] which have found applications in various physical problems, such as the description of the fractional statistics [2], the bosonization of supersymmetric quantum mechanics [3, 4] and in the n -particle integrable systems [4, 5]. Among these various deformations and extensions, we mention the generalized deformed oscillator algebras (GDOA's) [6] and the G -extended oscillator algebras [7] where G is some finite group, e.g in the case of Calogero-model $C_\lambda = \mathcal{Z}_\lambda$ is the cyclic group of order λ . This latter, have recently proved very useful in the context of supersymmetric quantum mechanics and some of its variants [7, 8], and also in the study of coherent states in nonlinear quantum optics [2].

In this paper, we recall some facts concerning the C_λ -extended oscillator algebras which were introduced as a generalization of Calogero-Vasiliev algebras [4, 9]. In the second section we give some of their mathematical properties, and we extend the commutation relation between annihilation and creation operators a and $a^\dagger$ to any indices n and m , as we will see for the positive indices it is straightforward. However, for the formal negative

¹Permanent Address: Université Mohammed V, Faculté des Sciences, Département de Physique, LPT, Av. Ibn Batouta, B.P. 1014, Agdal, Rabat, Morocco

²Junior Associate at The Abdus Salam ICTP, Trieste, Italy.
Permanent Address: Université Moulay Ismail, Faculté des Sciences et Techniques, Département de Mathématiques, Laboratoire de Physique Mathématique, Boutalamine B.P.509, Errachidia, Morocco.

powers one consider the Bargmann representation [9], indeed in such representation the generators a and $a^\dagger$ are identified with differential operators acting on the Bargmann space. Section three is devoted to the construction of w_∞ -algebra generators in terms of the pair a and $a^\dagger$ of the C_λ -extended oscillator algebras, we begin by the case of the Virasoro generators. We enclose by a conclusion and remarks.

2 Review on C_λ -extended oscillator algebras

As a generalization of the Calogero-Vasiliev algebras, the C_λ -extended oscillator algebras (also called generalized deformed oscillator algebras (GDOA's)), denoted A^λ , $\lambda = 2, 3, \dots$, are defined by

$$\begin{aligned} [N, a^\dagger] &= a^\dagger, & [N, P_\mu] &= 0, \\ [a, a^\dagger] &= I + \sum_{\mu=0}^{\lambda-1} \alpha_\mu P_\mu, & a^\dagger P_\mu &= P_{\mu+1} a^\dagger, \end{aligned} \quad (1)$$

together with their hermitian conjugates, and

$$\begin{aligned} P_\mu &= \frac{1}{\lambda} \sum_{\nu=0}^{\lambda-1} e^{\frac{2\pi i \nu(N-\mu)}{\lambda}}, & \sum_{\mu=0}^{\lambda-1} P_\mu &= 1 & P_\mu P_\nu &= \delta_{\mu,\nu} P_\nu, \\ \sum_{\mu=0}^{\lambda-1} \alpha_\mu &= 0, & \sum_{\nu=0}^{\mu-1} \alpha_\nu &> -1, & \mu &= 1, \dots, \lambda-1. \end{aligned} \quad (2)$$

where $\alpha_\mu \in \mathbf{R}$, N is the number operator and P_μ are the projection operators on subspaces $F_\mu = \{|k\lambda - \mu\rangle | k = 0, 1, 2, \dots\}$ of the Fock space F which is portioning into λ subspaces.

The operators a and $a^\dagger$ are defined by

$$a^\dagger a = F(N), \quad a a^\dagger = F(N+1), \quad (3)$$

where $F(N) = N + \sum_{\mu=0}^{\lambda-1} \beta_\mu P_\mu$, $\beta_\mu = \sum_{\nu=0}^{\mu-1} \alpha_\nu$, which is a fundamental concept of deformed oscillators. Let's denote the basis states of subspaces F_μ by $|n\rangle = |k\lambda + \mu\rangle \simeq (a^\dagger)^n |0\rangle$ where $a|0\rangle = 0$, $|0\rangle$ is the vacuum state. The operators a , $a^\dagger$ and N act on F_μ as follows

$$N|n\rangle = n|n\rangle, \quad a^\dagger|n\rangle = \sqrt{F(N+1)}|n+1\rangle, \quad a|n\rangle = \sqrt{F(N)}|n-1\rangle. \quad (4)$$

According to these relations, a and $a^\dagger$ are the annihilation and the creation operators respectively.

Particularly, if $\lambda = 2$, we have two projection operators $P_0 = \frac{1}{2}(I + (-1)^N)$ and $P_1 = \frac{1}{2}(I - (-1)^N)$ on the even and odd subspaces of the Fock space F , and the relations of (1) are restricted to

$$[N, a^\dagger] = a^\dagger, \quad [a, a^\dagger] = I + \kappa K, \quad \{K, a^\dagger\} = 0, \quad (5)$$

with their hermitian conjugates, where $K = (-1)^N$ is the Klein operator and κ is a real parameter. These relations define the so-called Calogero-Vasiliev algebra.

The C_λ -extended oscillator algebras are seen as deformation of G -extended oscillator algebras, where G is some finite group, appeared in connection with n -particle integrable models. In the former case, G is the symmetric group S_n . So, for two particles S_2 , can be realized in terms of K and S_2 -extended oscillator algebra becomes a generalized deformed oscillator algebra (GDOA) also known as the Calogero-Vasiliev or modified oscillator algebra. In the C_λ -extended oscillator algebras, $G \equiv C_\lambda$ is the cyclic group of order λ , $C_\lambda = \{1, K, \dots, K^{\lambda-1}\}$. So, these algebras have a rich structure since they depend upon λ independent real parameters, $\alpha_0, \alpha_1, \dots, \alpha_{\lambda-1}$.

Thus, these algebras can be also rewritten in terms of Klein operator, and instead of (1) we have

$$[N, a^\dagger] = a^\dagger, \quad [N, K] = 0, \quad K^\lambda = I, \quad (6)$$

$$[a, a^\dagger] = I + \sum_{r=1}^{\lambda-1} \kappa_r K^r, \quad a^\dagger K = e^{\frac{-2\pi i}{\lambda}} K a^\dagger, \quad (7)$$

together with their hermitian conjugates, where $\kappa_\mu \in \mathbf{C}$ and $\kappa_\mu^* = \kappa_{\lambda-\mu}$.

To see the equivalence between (1) and (6-7), we note that C_λ has λ inequivalent one-dimensional unitary irreducible matrix representations Γ^μ , $\mu = 0, 1, \dots, \lambda - 1$. The projection operator on the carrier space of Γ^μ can be written as

$$P_\mu = \frac{1}{\lambda} \sum_{\nu=0}^{\lambda-1} (\Gamma^\mu(K^\nu))^* K^\nu = \frac{1}{\lambda} \sum_{\nu=0}^{\lambda-1} e^{-2i\pi\mu\nu} K^\nu, \quad (8)$$

and α_μ can be given in terms of κ_μ as follows

$$\alpha_\mu = \sum_{\nu=0}^{\lambda-1} e^{\frac{-2i\pi\mu\nu}{\lambda}} \kappa_\nu, \quad \mu = 0, 1, \dots, \lambda - 1. \quad (9)$$

The C_λ -extended oscillator Hamiltonian defined in the bosonic Fock space representation as usual by

$$H_0 = \frac{1}{2} \{a, a^\dagger\}. \quad (10)$$

Owing to the relations of (6-7) the Hamiltonian can be rewritten in terms of the projection operators as

$$H_0 = N + \frac{1}{2}I + \sum_{\mu=0}^{\lambda-1} \gamma_\mu P_\mu, \quad \gamma_\mu \equiv \frac{1}{2}(\beta_\mu + \beta_{\mu+1}), \quad \beta_\mu = \sum_{\nu=0}^{\mu-1} \alpha_\nu. \quad (11)$$

Its eigenstates are $|n\rangle = |k\lambda + \mu\rangle$ and their eigenvalues are given by

$$E_{k\lambda+\mu} = k\lambda + \mu + \gamma_\mu + \frac{1}{2}, \quad k = 0, 1, \dots, \quad \mu = 0, 1, \dots, \lambda - 1. \quad (12)$$

In each subspace F_μ , the spectrum of H_0 is harmonic, but the λ infinite sets of equally spaced energy levels, corresponding to $\mu = 0, 1, \dots, \lambda - 1$, can be shifted with respect to each other by some amounts depending upon the parameters $\alpha_0, \alpha_1, \dots, \alpha_{\lambda-1}$.

In what follows, we will extend the commutation relations between $a^\dagger$ and a . First for the positive indices the extension is straightforward then from the following commutation relation between the generators $a^\dagger$, a and K .

$$[a, a^\dagger] = I + \sum_{r=1}^{\lambda-1} \kappa_r K^r, \quad a^\dagger K = e^{-2\pi i/\lambda} K a^\dagger, \quad (13)$$

one obtain easily after algebraic manipulation the following relations [17]

$$[a, (a^\dagger)^m] = (m + \sum_{r=1}^{\lambda-1} f_r \kappa_r K^r) (a^\dagger)^{m-1}, \quad (14)$$

where the functions f_r are given by

$$f_r = \sum_{p=0}^{m-1} e^{-i \frac{2r\pi}{\lambda} p}, \quad r = 2, 3, \dots, \lambda - 1, \quad f_1 = 1. \quad (15)$$

The negative values $m < -1$ are described as monomials of $a^\dagger$ with negative powers, which act as differential operators on the Bargmann representation [9]. For any order n of a one obtain,

$$[a^n, (a^\dagger)^m] = \sum_{\alpha=0}^{n-1} \sum_{l=0}^{\alpha} (m + F x^\alpha) \beta_l (a^\dagger)^{m-l-1} a^{n-l-1}, \quad (16)$$

where

$$F = \sum_{r=1}^{\lambda-1} f_r \kappa_r K^r, \quad x = e^{\frac{-2\pi i}{\lambda}}, \quad (17)$$

and such that

$$\begin{aligned} a^n (a^\dagger)^{m-1} &= \beta_n (a^\dagger)^{m-n-1} + \beta_{n-1} (a^\dagger)^{m-n} a + \beta_{n-2} (a^\dagger)^{m-n+1} a^2 + \dots + \\ &\beta_3 (a^\dagger)^{m-4} a^{n-3} + \beta_2 (a^\dagger)^{m-3} a^{n-2} + \beta_1 (a^\dagger)^{m-2} a^{n-1} + \beta_0 (a^\dagger)^{m-1} a^n. \end{aligned} \quad (18)$$

Owing to the relations (13) and (14) the parameters β_i , $i = 0, 1, 2, \dots, n - k, \dots, n$, are expressed by the following equalities

$$\begin{aligned}
\beta_n &= \prod_{i=1}^n \left((m-i) + Fx^{n-i} \right). \\
\beta_{n-1} &= \prod_{i=1}^{n-1} \left((m-i) + Fx^{n-i} \right) + \sum_{\mu=2}^{n-2} \prod_{i=1}^{n-\mu} \left((m-i) + Fx^{n-i} \right) \prod_{i=n-(\mu-1)}^{n-1} \left((m-i) + Fx^{\mu-i-1} \right) + \\
&\quad \left(2(m-1) + \sum_{j=n-2}^{n-1} Fx^j \right) \prod_{i=2}^{n-1} \left((m-i) + Fx^{n-i-1} \right). \\
&\quad \cdot \\
&\quad \cdot \\
&\quad \cdot \\
\beta_{n-k} &= \prod_{i=1}^{n-k} \left((m-i) + Fx^{n-i} \right) + \prod_{i=1}^{n-k-1} \left((m-i) + Fx^{n-i} \right) \left(k(m-(n-k)) + \sum_{j=0}^{k-1} Fx^j \right) + \\
&\quad \sum_{\alpha=k+2}^{n-2} \prod_{i=1}^{n-\alpha} \left((m-i) + Fx^{n-i} \right) \left[\prod_{i=n-\alpha+1}^{n-k} \left((m-i) + Fx^{n-i-1} \right) + \right. \\
&\quad \left. \sum_{\mu=n-\alpha+1}^{n-k-1} \prod_{i=n-\alpha-1}^{n-\mu} \left((m-i) + Fx^{n-i-1} \right) \prod_{i=n-\mu+1}^{n-k} \left((m-i) + Fx^{\mu-i-2} \right) + \right. \\
&\quad \left. \sum_{l=2}^k \left(l(m-(n-\alpha+1)) + \sum_{j=\alpha-l-1}^{n-4} Fx^j \right) \prod_{i=n-\alpha+2}^{n-l} \left((m-i) + Fx^{n-i-l} \right) \right] + \\
&\quad \sum_{l=2}^k \left(l(m-1) + \sum_{j=n-2}^{n-1} Fx^j \right) \left[\prod_{i=2}^{n-k} \left((m-i) + Fx^{n-i-1} \right) + \right. \\
&\quad \left. \sum_{\mu=k+1}^{n-4} \prod_{i=2}^{n-\mu} \left((m-i) + Fx^{n-i-1} \right) \prod_{i=n-\mu+1}^{n-k-2} \left((m-i) + Fx^{\mu-i-2} \right) + \right. \\
&\quad \left. \left((m-2) + Fx^{n-3} \right) \left((k+2-l)(m-3) + \sum_{j=n-5}^{n-4} Fx^j \right) \prod_{i=n-k-1}^{n-k} \left((m-i) + Fx^{n-i-k} \right) \right] + \\
&\quad \left((k+1)(m-1) + \sum_{j=2}^{n-1} Fx^j \right) \prod_{i=2}^{n-2} \left((m-i) + Fx^{n-i-2} \right), \quad k = 2, \dots, n-4 \\
&\quad \cdot \\
&\quad \cdot \\
&\quad \cdot \\
\beta_3 &= \prod_{i=1}^3 \left((m-i) + Fx^{n-i} \right) + \prod_{i=1}^2 \left((m-i) + Fx^{n-i} \right) \left((n-3)(m-3) + \sum_{j=0}^{n-4} Fx^j \right) + \\
&\quad \sum_{\mu=2}^{n-2} \left(\mu(m-1) + \sum_{j=n-\mu}^{n-1} Fx^j \right) \left((m-2) + Fx^{n-\mu-1} \right) \left((n-\mu-1)(m-3) + \sum_{j=0}^{n-\mu-2} Fx^j \right) \\
\beta_2 &= \prod_{i=1}^2 \left((m-i) + Fx^{n-i} \right) + \sum_{\mu=2}^{n-1} \left(\mu(m-1) + \sum_{j=n-\mu}^{n-1} Fx^j \right) \left((m-2) + Fx^{n-\mu-1} \right) \\
\beta_1 &= \left(n(m-1) + \sum_{j=0}^{n-1} Fx^j \right) \\
\beta_0 &= 1
\end{aligned}$$

3 Modified w_∞ -symmetry

In this section, we introduce the new modified w_∞ -symmetry using the C_λ -extended oscillator generators. Here we will adopt the approach for deformed or undeformed case where the elements of the w_∞ -algebra ω_m^s are expressed in terms of the pair $a^\dagger$ and a of quantum oscillator with or without deformation. Before going on, let us first restrict ourselves to the case of classical Virasoro algebra generated by the operators $\omega_m^2 = \ell_m$

3.1 Classical conformal symmetry and its representations

For reader convenience we remind some information about the conformal symmetry termed also Virasoro symmetry which was first introduced in the context of string theories and it is relevant to any theory in (2+1)-dimensional space-time which possesses conformal invariance. The conformal symmetry consists of all general transformations

$$z \rightarrow z + \epsilon(z), \quad \bar{z} \rightarrow \bar{z} + \bar{\epsilon}(\bar{z}) \quad (20)$$

where $\epsilon(z)$ and $(\bar{\epsilon}(\bar{z}))$ are an infinitesimal analytical (anti-analytical) functions. It can be represented as an infinite Laurent series

$$\epsilon(z) = \sum_n \epsilon_n z^{n+1}, \quad n \in \mathbb{Z}, \quad (21)$$

and an analogous formula holds for $\bar{\epsilon}(\bar{z})$.

These mapping are generated by the differential operators

$$\ell_m = -z^{m+1} \partial_z \quad \text{and} \quad \bar{\ell}_m = -\bar{z}^{m+1} \partial_{\bar{z}}. \quad (22)$$

The operators ℓ_m satisfy the commutator relations

$$[\ell_m, \ell_n] = (m - n) \ell_{m+n}, \quad (23)$$

and an analogous formula holds for the $\bar{\ell}_m$. Hence, this means that both the ℓ_m and the $\bar{\ell}_m$ span an infinite-dimensional Lie algebra; moreover, these two algebras are combined as a direct sum, $[\ell_m, \bar{\ell}_m] = 0$. The algebra defined by (23) is known as the classical conformal or Virasoro algebra. We shall denote that the above algebra admits a unique 1-dimensional central extension

$$\mathcal{L}_c = \mathcal{L} \oplus \mathcal{C}c, \quad (24)$$

with the following commutation relations

$$[\ell_m, c] = 0, \quad [\ell_m, \ell_n] = (m - n) \ell_{m+n} + c \frac{(m^3 - m)}{12} \delta_{m+n, 0}, \quad (25)$$

where the value of the central charge c is the parameter of the theory in the quantum field theory context (e.g $c = 1$ for the free boson). The unitary representation of the above algebra are well known (see e.g [10]). In what follows, we give some representations discussed in the literature,

(i) the oscillatory representation of the elements ℓ_m

$$\ell_m = (a^\dagger)^{m+1}a,$$

(ii) the Sugawara construction, where the generators ℓ_m are constructed in terms of infinite set of free boson operators ($b_m, b_m^\dagger = b_{-m}, m = 1, 2, \dots$) as

$$\ell_m = \frac{1}{2} \sum_{j \in \mathbb{Z}} : b_{m-j} b_j :,$$

where the colons indicate normal ordering,

(iii) the infinite matrix realization

$$\ell_m = \sum_{j \in \mathbb{Z}} E_{m, m-j},$$

where $E_{k,j}$ are the infinite dimensional matrix with value 1 at the (k, j) entry and 0 at all other entries [10].

Here we consider the simplest one which is given by oscillatory representation. In fact using the commutation relations of the quantum mechanics, we obtain the following commutator

$$[a, (a^\dagger)^m] = m(a^\dagger)^{m-1}. \quad (26)$$

From this commutator, one see easily that the Virasoro operators for the unidimensional harmonic oscillators can be defined by

$$\ell_m = (a^\dagger)^{m+1}a, \quad (27)$$

which satisfy the following centerless Virasoro algebra

$$[\ell_m, \ell_n] = (m - n)\ell_{m+n}. \quad (28)$$

3.2 C_λ -deformed Virasoro algebra

First, we recall that the deformation of the conformal algebra was first introduced by Curtright and Zachos [11] and investigated on many occasions by many authors [12, 13, 14], and defined by the following q -commutation relation

$$[\ell_m, \ell_n]_q = q^{m-n}\ell_m\ell_n - q^{n-m}\ell_n\ell_m = \frac{(q^{m-n} - q^{n-m})}{q - q^{-1}}\ell_{m+n}. \quad (29)$$

Turn now to the construction of the C_λ -deformed Virasoro algebra. To do this, we will adopt the approach for undeformed case [15] where the generators ℓ_m are constructed from one undeformed oscillator pair $a^\dagger$ and a eq (27) as the infinite-dimensional extension of the following realization of $sp(2) \sim o(2, 1)$

$$\ell_{-1} = a, \quad \ell_0 = (a^\dagger)a, \quad \ell_1 = (a^\dagger)^2a. \quad (30)$$

The extension to positive indices m is straightforward

$$\ell_m(\kappa) = (a^\dagger)^{m+1}a, \quad (31)$$

and for the negative values ($m < -1$, which makes this construction formal) the generators $\ell_m(\kappa)$ are described by the nonanalytic dependence (monomials of $a^\dagger$ with negative powers), which act as the differential operators in the Bargmann representation [9].

Then from the previous equations (14, 15), one gets the commutation relations for the generators $\ell_m(\kappa)$ and the Klein operator K [17]

$$[\ell_m(\kappa), \ell_n(\kappa)] = (m - n)\ell_{m+n}(\kappa) + \sum_{r=1}^{\lambda-1} (e^{(n+1)r(2\pi i/\lambda)} - e^{(m+1)r(2\pi i/\lambda)})\kappa_r K^r \ell_{m+n}(\kappa) \quad (32)$$

$$[\ell_m(\kappa), K] = g^m(\lambda)\ell_m(\kappa)K, \quad (33)$$

where the functions $g^m(\lambda)$ are given by

$$g^m(\lambda) = (1 - e^{\frac{2\pi i(m+1)}{\lambda}}). \quad (34)$$

From the above expression of the function $g^m(\lambda)$, one note that the generators $\ell_m(\kappa)$ commutes with the Klein-operator only in the case where $(1 - e^{\frac{2\pi i(m+1)}{\lambda}} = 0)$ that is when $m+1 = k\lambda$, $k \in \mathcal{Z}$. Note also that the ordinary Virasoro algebra is obtained for $\kappa_r \rightarrow 0$. However, when $\kappa_r \neq 0$ is something new. In the case of the Calogero-Vasiliev case, $\lambda = 2$ ($r = 1$), the relation (14) becomes

$$[a, (a^\dagger)^m] = (m + (1 - (-1)^m)\kappa_1 K)(a^\dagger)^{m-1}, \quad (35)$$

In what follows, we investigate the possible cases of the indices m : (I) $m = 2k$ ($k \in \mathcal{Z}$) even and (II) $m = 2k + 1$ odd. In (I) the above commutation relation becomes

$$[a, (a^\dagger)^{2k}] = 2k(a^\dagger)^{2k-1}, \quad (36)$$

and the obtained C_2 -Virasoro algebra is given by

$$[\ell_{2k}(\kappa), \ell_{2l}(\kappa)] = (2k - 2l)\ell_{2k+2l}, \quad [\ell_{2k}, K] = 2\ell_{2k}K. \quad (37)$$

In the case (II), we have

$$[a, (a^\dagger)^{2k+1}] = (2k + 1 + 2\kappa_1 K)(a^\dagger)^{2k}, \quad (38)$$

we obtain the following algebra

$$[\ell_{2k+1}(\kappa), \ell_{2l+1}(\kappa)] = (2k - 2l)\ell_{2k+2l+2}, \quad [\ell_{2k+1}, K] = 0. \quad (39)$$

Then, the conformal algebra is not changed when the two indices (m, n) are both even or odd, however when one indices is even and the other is odd the obtained Virasoro algebra is changed, we have

$$[\ell_{2n}(\kappa), \ell_{2m+1}(\kappa)] = 2(m - n)\ell_{2m+2n+1}(\kappa) + (1 - 2\kappa_1 K)\ell_{2m+2n+1}, \quad (40)$$

$$[\ell_{2n}(\kappa), K] = 2\ell_{2n}(\kappa)K \quad (41)$$

$$[\ell_{2m+1}(\kappa), K] = 0 \quad (42)$$

3.3 C_λ -deformed w_∞ -algebra

In this section, we will extend the construction done in the previous section on w_∞ -algebra which is seen as a generalization of Virasoro algebra. This algebra is interesting in the conformal field theory point of view because it contains the centerless Virasoro algebra as subalgebra and thus may be viewed as an extended conformal symmetry with an infinite set of additional symmetries. The w_∞ -algebra contains generators ω_m^s of spin $2, 3, 4, \dots$. The defining commutation relations are

$$[\omega_m^s, \omega_n^t] = ((t-1)m - (s-1)n)\omega_{m+n}^{s+t-2}. \quad (43)$$

The generators ω_m^2 generate a classical Virasoro subalgebra. The conformal properties of these generators can be viewed from the commutation relation

$$[\omega_m^2, \omega_n^t] = ((t-1)m - n)\omega_{m+n}^t. \quad (44)$$

which implies that ω_m^t can be regarded as the algebra Fourier modes of primary fields with the conform spin t . On the other hand w_∞ -algebra can have a natural geometric interpretation as area preserving diffeomorphisms of 2-manifolds e.g on the two-dimensional cylinder $\mathcal{R} \times S^1$ with canonical coordinates x and y it can be represented by Poisson bracket of the smooth function $\omega_m^s = e^{mx}y^{s-1}$. However, the standard central extension of Virasoro algebra can not be extended to the full algebra w_∞ , which should therefore be viewed as a classical W -algebra. The W_∞ -algebra, which was first introduced in refs [18], is a deformation of w_∞ which is such that the standard central term in the Virasoro subalgebra can be extended to the whole algebra. W_∞ -algebra can thus be viewed as the quantum version of w_∞ . This has been nicely illustrated in the context of W gravity. Denote the generators of W_∞ -algebra of spin s by V_m^i , where $s = i + 2$, and the index i ranges from 0 to ∞ . The defining commutation relations of W_∞ -algebra can be written as

$$[V_m^i, V_n^j] = \sum_{l>0} g_{2l}^{ij}(m, n)V_{m+n}^{i+j-2l} + c_i(m)\delta^{ij}\delta_{m+n}. \quad (45)$$

The structure constant $g_{2l}^{ij}(m, n)$ and the central terms $c_i(m)$ are completely fixed by the Jacobi identities and take the following form

$$c_i(m) = m(m^2 - 1)(m^2 - 4)\dots(m^2 - (i+1)^2)c_i, \quad (46)$$

where the central charges c_i are given by

$$c_i = \frac{2^{2i-3}i!(i+1)!}{(2i+1)!!(2i+3)!!}. \quad (47)$$

The structure constant $g_{2l}^{ij}(m, n)$ are given by

$$g_{2l}^{ij}(m, n) = \frac{1}{2(l+1)}\phi_l^{ij}N_l^{ij}(m, n), \quad (48)$$

where the $N_l^{ij}(m, n)$ are expressed as

$$N_l^{ij}(m, n) = \sum_{k=0}^{l+1} (-1)^k \binom{l+1}{k} [i+1+m]_{(l+1-k)} [i+1+m]_k [j+1+n]_k [j+1-n]_{(l+1-k)} \quad (49)$$

where $[x]_n = \frac{\Gamma(x+1)}{\Gamma(x+1-n)}$. Finally, the ϕ_l^{ij} are given by

$$\phi_l^{ij} = \sum_{k>0} \frac{\left(\frac{-1}{2}\right)_k \left(\frac{3}{2}\right)_k \left(\frac{-1l}{2} - \frac{-1}{2}\right)_k \left(\frac{-1l}{2}\right)_k}{k! \left(-i - \frac{-1}{2}\right)_k \left(-j - \frac{-1}{2}\right)_k \left(i + j - l + \frac{5}{2}\right)_k}, \quad (50)$$

with the notation $(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)} = x(x+1)\dots(x+n-1)$.

Now, turn the realization of w_∞ -algebra generators in terms of the pair $a^\dagger$ and a of the C_λ -extended oscillator algebras by extending the realization (31). To begin, let's define the following operators

$$\omega_m^s = (a^\dagger)^s a^m \quad (51)$$

By using the previous tools (16-19) the commutation relations of the operators ω_m^s are

$$[\omega_m^s, \omega_n^t] = (\Xi^{(s,m)} - \Xi^{(t,n)}) \omega_{m+n-(l+1)}^{s+t-(l+1)}, \quad (52)$$

where

$$\Xi^{(s,m)} = m(t+Fx^s)\beta_0^{[s]} + (m-1)(t+Fx^{1+s})\beta_1^{[s]} + (m-2)(t+Fx^{2+s})\beta_2^{[s]} + \dots + (t+Fx^{(m-1)+s})\beta_{m-1}^{[s]}, \quad (53)$$

and

$$\Xi^{(t,n)} = n(s+Fx^t)\beta_0^{[t]} + (n-1)(s+Fx^{1+t})\beta_1^{[t]} + (n-2)(s+Fx^{2+t})\beta_2^{[t]} + \dots + (s+Fx^{(n-1)+t})\beta_{n-1}^{[t]}, \quad (54)$$

where the notation $\beta_l^{[n]}$ means that each x^j in β_l becomes x^{jn} and β_l are given by eq (19). Owing to the commutation relations of (13) and the definition of ω_m^s , we obtain the following commutation relations between the K (Klein operator) and the generators ω_m^s .

$$[\omega_m^s, K] = (x^{s+m} - 1)K\omega_m^s. \quad (55)$$

We note that only in the case where $s + m = \lambda$, the generators ω_m^s commute with the Klein operator.

In what follows, we will examine some particular cases obtained as limits. First, setting $m = 1$ and $s \rightarrow s + 1$, we recover the C_λ -extended Virasoro algebra (32). Second, the C_λ -extended $sp(2)$ is generated by ω_1^0, ω_1^2 and ω_1^1 which satisfy

$$[\omega_1^0, \omega_1^1] \equiv \sum_{\alpha, l} \Phi_{\alpha, l}^{[0, 1, 1, 1]} \omega_{1-l}^{-l} \quad (56)$$

$$[\omega_1^2, \omega_1^1] \equiv \sum_{\alpha, l} \Phi_{\alpha, l}^{[2, 1, 1, 1]} \omega_{1-l}^{2-l} \quad (57)$$

$$[\omega_1^2, \omega_1^0] \equiv \sum_{\alpha, l} \Phi_{\alpha, l}^{[2, 1, 0, 1]} \omega_{1-l}^{1-l}, \quad (58)$$

where

$$\sum_{\alpha, l} \Phi_{\alpha, l}^{[0, 1, 1, 1]} \omega_{1-l}^{-l} = \left[\sum_{r=1}^{\lambda-1} \kappa_r K^r (1-x) + 1 \right] \omega_1^0 \quad (59)$$

$$\sum_{\alpha,l} \Phi_{\alpha,l}^{[2,1,1,1]} \omega_{1-l}^{2-l} = \left[\sum_{r=1}^{\lambda-1} \kappa_r K^r (1+x^r) x(x-1) - 1 \right] \omega_1^2 \quad (60)$$

$$\sum_{\alpha,l} \Phi_{\alpha,l}^{[2,1,0,1]} \omega_{1-l}^{1-l} = \left[\sum_{r=1}^{\lambda-1} \kappa_r K^r (1+x^r) (x^2-1) - 2 \right] \omega_1^1, \quad (61)$$

with x is given by (17). The corresponding C_λ -extended Casimir operator is

$$C(\lambda) = (\omega_1^1)^2 - \frac{1}{2} \{ \omega_1^2, \omega_1^0 \}, \quad (62)$$

evidently the C_λ -extended Casimir don't commute with all generators (53-55), for example

$$\begin{aligned} [C(\lambda), \omega_1^1] &= -\frac{1}{2} [\{ \omega_1^2, \omega_1^0 \}, \omega_1^1] \\ &= \frac{1}{2} \left[\sum_{r=1}^{\lambda-1} \kappa_r K^r (x^{3r} - (1+x^r)x - (x^r + x^{2r})x + 1)(x-1) \right] \omega_2^2 \\ &\quad - \left[\sum_{r=1}^{\lambda-1} \kappa_r K^r (x^r + x^{2r})x - 1 \right] \left[1 + \frac{1}{2} \sum_{r=1}^{\lambda-1} \kappa_r K^r (1+x^r) \right] (x-1) \omega_1^1 \end{aligned} \quad (63)$$

4 Conclusion and remarks

In the present paper, we have found the new deformed w_∞ -symmetry by adopting the approach for undeformed case where the elements of the w_∞ -algebras ω_m^s may be expressed via the pair $a^\dagger$ and a of quantum oscillator. We have shown that in the Calogero-Vasiliev case $\lambda = 2$, only in the case associated with odd and even indices the obtained symmetries are changed. Finally, note that in the same way one can construct the C_λ -extended oscillator algebras of the supersymmetric extension of w_∞ -algebra by using the super-realization of C_λ -extended oscillator algebras.

Acknowledgements

J. Douari acknowledges the hospitality of the IHES, France. E. H. El Kinani is grateful to the Abdus-Salam ICPT, Trieste, Italy for its hospitality.

References

- [1] V. G. Drinfeld, in: Proc. Int. Congress of Mathematicians, Berkeley, CA, 1986. ed. A. M. Cleason (AMS, Providence, RI, 1987) p. 798; M. Jimbo, Lett. Math. Phys. 10 (1985) 63; 11 (1986) 247.
- [2] J. M. Leinaas and J. Myrheim, Nuovo Cimento B37 (1977) 1; F. Wilczek, Fractional statistics and anyon superconductivity (World scientific, singapore, 1990); M. Daoud and J. Douari, Mod. Phys. Lett. A18 (2003) 913; O. W. Greenberg, Phys. Rev. Lett. 64 (1990) 705; Phys. Rev. D43 (1991) 4111; S. Chaturvedi and V. Srinivasan, Phys. Rev. A44 (1991) 8024.

- [3] M. S. Plyushchay, Ann. Phys. (NY) 245 (1996) 339; D. Bonatsos and C. Daskaloyannis, Phys. Lett. B307 (1993) 100.
- [4] T. Brzezński, I. L. Egusquiza and A. J. Macfarlane, Phys. Lett. B311 (1993) 202.
- [5] M. A. Vasiliev, Int. J. Mod. Phys. A6 (1991) 1115; A. P. Polychronakos, Phys. rev. lett. 69 (1992) 703; L. Brink, T. H. Hansson and M. A. Vasiliev, Phys. lett. B286 (1992) 109.
- [6] L. C. Biedenhorn, J. Phys. A22 (1989) L873; A. J. Macfarlane, J. Phys. A22 (1982) 4581; A. Junussis, J. Phys. A26 (1982) L233; C. Quesne, N. Vansteenkiste, J. Phys. A28 (1995) 7019; Helv. Phys. Acta 69 (1996) 141.
- [7] C. Quesne, N. Vansteenkiste, Phys. lett. A240 (1998) 21.
- [8] C. Quesne, N. Vansteenkiste, Helv. Phys. Acta 72 (1999) 71.
- [9] C. Quesne, Phys. lett. A272 (2000) 313, Ann. Phys. 293(2001)147.
- [10] V.G.Kac and A.B Raina, Bonbay Lectures on the Highest Weight Representations of Infinite dimensional Lie Algebras (World Scietific, Singapor, 1987).
- [11] T.Curtright and C.Zachos, Phys.lett.B 243(1990)237.
- [12] M.Chaichan, D.E.Ilinas and Z.Popowicz, Phys.lett.B 248(1990) 95.
- [13] N.Aizawa and H.Sato, Phys.lett.B 256(1991) 185.
- [14] E.H.El Kinani and M.Zakkari, Phys.lett.B 357(1995) 105.
- [15] M.Chaichan, P.Kulish and J.Lukierski,Phys.lett.B 237(1990) 401.
- [16] M. A. Vasiliev, Int. J. Mod. Phys. A6 (1991) 1115.
- [17] E. H. El Kinani, Phys. lett. A 296 (2002) 259.
- [18] C.N.Pope, L.J Romans and X.Shen, Phys. lett. B 236 (1990) 137.